arXiv:0902.0446v1 [nucl-th] 3 Feb 2009

# Transverse momentum distribution of muon pairs from relativistic heavy ion collisions

Jan-e Alam[1], Tetsufumi Hirano[2] Jajati K Nayak[1] and Bikash Sinha[1]
*Variable Energy Cyclotron Centre, 1/AF, Bidhan Nagar, Kolkata 700 064, India and*
*Physics Department, The University of Tokyo, Tokyo 113-0033, Japan*

(Dated: February 3, 2009)

Recently NA60 collaboration has extracted the inverse slope parameters from the dimuon spectra originating from In+In collisions at $\sqrt{s_{\rm NN}} = 17.3$ GeV for various invariant mass region. They have observed that the inverse slope parameter as a function of invariant mass of the lepton pair drops beyond the $\rho$-peak. In the present work we study the slope parameters for the low mass region (LMR) and show that the experimentally observed behavior can be explained by assuming the formation of a partonic phase which reverts to hadronic phase through a very weak first order (or continuous) phase transition at a temperature $T_c \sim 175$ MeV. A strong first order transition fails to explain the data. It is observed that a scenario without the formation of a partonic phase does not reproduce the non-monotonic behaviour of the inverse slope parameter non typical of radial flow. It is argued that the variation of $T_{\rm eff}$ with invariant mass $M$ may be used to constrain the transition temperature.

PACS numbers: 25.75.-q,25.75.Dw,24.85.+p

Low mass lepton pairs are considered as useful tools to probe the thermodynamic state of the matter presumably formed in nuclear collisions at relativistic energies [1, 2, 3, 4, 5, 6]. Unlike hadrons which bring the information of the state of the matter when they cease to interact - the lepton pairs probe the entire stage of the evolution history - from the initial formation time to the final freeze-out time of the fireball. Recently high quality data on dimuon spectra from NA60 collaboration [7] has been made available for both the kinematical variables - the transverse momentum ($p_T$) and the invariant mass ($M$) of the pairs. They observed a non-monotonic behaviour of the inverse slope parameter, $T_{\rm eff}$, extracted from the transverse mass spectra of the pairs - as a function of invariant mass and indicated that such a trend may be originated due to the radiation of lepton pairs from a partonic phase formed initially in the collisions.

The probability that a muon pair of invariant mass, $M$ and transverse momentum, $p_T$ will emit from a thermal system at temperature $T$ is determined by the Boltzmann factor $\sim e^{-\sqrt{M^2+p_T^2}/T}$. For a dynamically evolving system like the one produced after nuclear collisions at ultra-relativistic energies - the temperature decreases with time because a part of the thermal energy is spent to allow the collective motion of the system. Consequently the Boltzmann factor is modified to $\sim e^{-\sqrt{M^2+p_T^2}/T_{\rm eff}}$, where $T_{\rm eff} \sim T_{\rm th} + M v_r^2$ - here the first term represent thermal part and the second term stands for the flow part for flow velocity $v_r$. It is expected that the large $M$ thermal pairs will be emitted from early time when temperature is large and flow velocity is small and the small $M$ pairs will originate from the late stage of the evolution when the temperature is low but flow velocity is large. Therefore, the variation of the $T_{\rm eff}$ with $M$ may be treated as a chronometer of the heavy ion collisions.

In the present work, therefore, we focus on the transverse mass distribution of the lepton pairs formed in In + In collisions at 158A GeV beam energy at CERN-SPS and subsequently extract $T_{\rm eff}$. We assume the following two scenarios for the collisions: (i)In+In → quark gluon plasma (QGP) → mixed phase of quarks and hadrons → hadronic phase and (ii) In+In → hadronic phase and check by comparing with data which is the possible scenario realized in these collisions. In contrast to earlier studies [8, 9, 10, 11] we focus on the sensitivities of the slopes on the equations of state and transition temperature, $T_c$.

The rate of thermal dilepton production per unit space-time volume per unit four momentum volume is given by[1, 2, 3]

$$\frac{dN}{d^4p d^4x} = \frac{\alpha}{12\pi^4 M^2} L(M^2) {\rm Im}\Pi^{{\rm R}\mu}_{\mu} f_{\rm BE} \qquad (1)$$

where $\alpha$ is the electromagnetic coupling, ${\rm Im}\Pi^{\mu}_{\mu}$ is the imaginary part of the retarded photon self energy and $f_{\rm BE}(E,T)$ is the thermal phase space factor for Bosons, where the energy $E$ should be replaced by $u^\mu p_\mu$ for an expanding system having four velocity $u^\mu$ at each space-time point of the system, $p^2 (= p_\mu p^\mu) = M^2$ is the invariant mass square of the lepton pair and

$$L(M^2) = \left(1 + 2\frac{m^2}{M^2}\right)\sqrt{1 - 4\frac{m^2}{M^2}} \qquad (2)$$

arises from the final state leptonic current involving Dirac spinors, $m$ in Eq. 2 is the muon mass.

The major sources of thermal dileptons are quark-anti-quark annihilation in the QGP phase and the decays of light vector mesons ($\rho$, $\omega$ and $\phi$) in the hadronic phase. The other non-thermal sources are: (i) Drell-Yan process originating from the interactions of quarks and anti-quarks of the colliding nuclei and the decays of various mesons (*e.g.* $\pi$,$\eta$, $\omega$, $\rho$, $\eta'$, $\phi$ etc) after the fireball freeze-out. As the non-thermal contributions have been subtracted from the data under consideration we concentrate only on the thermal emissions. To evaluate the thermal dilepton spectra the static rate of emission ($dR/d^2m_Td M^2dy$) has to be convoluted with the space-time dynamics governed by the relativistic hydrodynamics as follows:

$$\begin{aligned}\frac{dN_{\gamma^*}}{m_T dm_T} &= 2\pi \sum_{phases} \int \left(\frac{dR_{\gamma^*}}{d^2m_T dy dM^2}\right)_{phase} \\ &\quad \times dM^2 dy d^4x \qquad (3)\end{aligned}$$

$d^4x$ is the 4 dimensional volume element, $m_T$ is the transverse mass and $y$ is the rapidity. The limits for integration over invariant mass, $M$ can be fixed according to the experimental measurements. For the experimental data for $M_{\rm min} \leq M \leq M_{\rm max}$, the transverse mass $m_T$ is defined as $m_T = \sqrt{M^2 + p_T^2}$, where $M = (M_{\rm min} + (M_{\rm max})/2$.

The emission rate from the QGP has been obtained from the following processes $q\bar{q} \rightarrow l^+l^-$, $q\bar{q} \rightarrow gl^+l^-$ $gq(\bar{q}) \rightarrow q(\bar{q})l^+l^-$) (see Ref. [12, 13] for details).

The following parametrization [5, 14] has been used to evaluate the dilepton emission rates from light vector mesons ($\rho$, $\omega$ and $\phi$):

$$\begin{aligned}\frac{d^2R_{\gamma^*}}{dM^2d^2p_Tdy} &= \frac{\alpha^2}{2\pi^3}L(M^2)f_{BE} \\ &\quad [\frac{f_V^2 M\Gamma_V}{(M^2-m_V^2)^2+(M\Gamma_V)^2} \\ &\quad +\frac{1}{8\pi}\frac{1}{1+exp((w_0-M)/\delta)} \\ &\quad \times(1+\frac{\alpha_s}{\pi})]. \qquad (4)\end{aligned}$$

The parameterizations of the vector mesons spectral function are consistent with the experimental data from $e^+ e^- \rightarrow V(\rho, \omega$ or $\phi)$ processes [5, 6, 14]. $f_V$ is the coupling between the electromagnetic current and vector meson fields, $m_V$ and $\Gamma_V$ are the masses and widths of the vector mesons and $\omega_0$ is the continuum threshold above which the asymptotic freedom is restored. We have taken $\alpha_s = 0.3$, $\delta = 0.2GeV$, $\omega_0 = 1.3$ GeV for $\rho$ and $\omega$. For $\phi$ we have taken $\omega_0 = 1.5$ GeV and $\delta = 1.5$ Gev. The electromagnetic current in terms of $\rho$, $\omega$ and $\phi$ field can be expressed as $J_\mu = J_\mu^\rho + J_\mu^\omega/3 - J_\mu^\phi/3$. Therefore, the contributions from $\omega$ and $\phi$ will be down by a factor of 9 compared to $\rho$. As we have included the continuum in the vector mesons spectral functions four pion annihilation process [15] is not considered here to avoid over counting.

The space time evolution of the system has been studied using ideal relativistic hydrodynamics with longitudinal boost invariance [16] and cylindrical symmetry [17]. The initial temperature($T_i$) and thermalization time ($\tau_i$) are constrained by the following equation [18] for an isentropic expansion:

$$T_i^3\tau_i \approx \frac{2\pi^4}{45\xi(3)}\frac{1}{4a_{\rm eff}}\frac{1}{\pi R_A^2}\frac{dN}{dy}. \qquad (5)$$

where, $dN/dy$= hadron multiplicity, $R_A$ is the effective radius (evaluated by using the formula $R_A \sim N_{part}^{1/3}$) of the system, $\xi(3)$ is the Riemann zeta function and $a = \pi^2 g/90$ ($g = 37$ in the QGP), $N_F$=number of flavours. The initial radial velocity, $v_r(\tau_i, r)$ and energy density, $\epsilon(\tau_i, r)$ profiles are taken as:

$$v(\tau_i, r) = 0 \qquad (6)$$

and

$$\epsilon(\tau_i, r) = \frac{\epsilon_0}{e^{\frac{r-R_A}{\delta}}+1} \qquad (7)$$

where the surface thickness, $\delta = 0.5$ fm. So far there is no consensus on the value of $T_c$, it varies from 151 MeV [19] to 192 MeV [20]. In the present work we assume $T_c = 175$ MeV. Results for a higher $T_c$ will also be discussed. In a quark gluon plasma to hadronic transition scenario - we use the bag model EOS for the QGP phase and for the hadronic phase all the resonances with mass $\leq 2.5$ GeV have been considered [21]. The transition region has been parametrized as follows [23]:

$$s = f(T)s_q + (1 - f(T))s_h \qquad (8)$$

where $s_q$ ($s_h$) is the entropy density of the quark (hadronic) phase at $T_c$ and

$$f(T) = \frac{1}{2}(1 + \tanh(\frac{T - T_c}{\Gamma})) \qquad (9)$$

the value of the parameter $\Gamma$ can be varied to make the transition strong first order or continuous.

The ratios of various hadrons measured experimentally at different $\sqrt{s_{\rm NN}}$ indicate that the the system formed in heavy ion collisions chemically decouple at a temperature ($T_{\rm ch}$) which is higher than the temperature for kinetic freeze-out ($T_f$)determined by the transverse spectra of hadrons [24]. Therefore, the system remains out of chemical equilibrium from $T_{\rm ch}$ to $T_f$. The chemical non-equilibration affects the dilepton yields at two levels: a) the emission rate through the phase space factor and b) the



FIG. 1: In variant mass spectra of dilepton from quark gluon plasma and hadronic phase at $T = 0.2GeV$.



FIG. 2: The data is for $m_T - M$ spectra measured by NA60 collaboration for In+In collision for $M$ range indicated. The solid line is the theoretical result.

space-time evolution of the matter through the equation state. The value of the chemical potential and the its inclusion in the EOS has been taken in to account following [25]. It is expected that the chemical potentials do not change much for the inclusion of resonances above $\Delta$.

To evaluate the $m_T$ distribution of dilepton originating from the decays of vector mesons after freeze-out the Cooper-Frye formula [26] has been used. For a special case of unstable vector mesons we need to know the thermal phase space factor corresponding to an unstable Boson which is given by

$$f_{\rm unstable} = \frac{g}{(2\pi)^3}\,\frac{1}{\exp(\rm p_0)/T - 1}\rho(M) \qquad (10)$$



FIG. 3: Same as Fig. 2 for $0.4 \leq M$ (GeV)$\leq 0.6$

.



FIG. 4: Same as Fig. 2 for $0.6 \leq M$ (GeV)$\leq 0.9$

.

where $p_0 = \sqrt{p^2 + M^2}$, $g$ is the statistical degeneracy and $\rho(M)$ is the spectral function of the vector meson under consideration. For stable particle $\rho(M)$ reduces to a Dirac delta function and consequently the usual phase space factor for a stable particle is recovered. Therefore, the $m_T$ distribution of dimuons from vector meson decay after the freeze-out is given by



FIG. 5: Same as Fig. 2 for $1.0 \leq M$ (GeV)$\leq 1.4$.

$$
\begin{aligned}
\frac{dN_{\gamma^*}}{m_T dm_T} &= 2\pi \int dr \int d\eta \int d\phi \, r\tau \\
&\times \left( m_T \cosh(y-\eta) - \frac{\partial \tau}{\partial r} p_T \cos\phi \right) \\
&\times \rho(M) \Gamma_{V\to\mu^+\mu^-}/\Gamma_V^{\rm tot} f_{\rm unstable} \\
&dM^2 dy \qquad (11)
\end{aligned}
$$

where $\Gamma_V^{\rm tot}$ is the total decay width of the vector meson, $V$. The contributions from Eqs. 3 and 11 are added for the description of the $m_T$ spectra of the data.

Before discussing the $m_T$ spectra it is important to mention here that initially there was no collective flow ($v(\tau_i, r) = 0$), the entire energy of the system was thermal. With the progress of time some part of the thermal energy get converted to the collective (flow) energy for a system undergoing hydrodynamic expansion. The measured $m_T$ spectra of muon pairs therefore, contains contributions from both thermal as well as collective degrees of freedom *i.e.* the inverse slope parameter, $T_{\rm eff}$ can be written as $T_{\rm eff} = T + Mv_r^2$ as mentioned earlier. Therefore, it is important to know the domains of $M$ where the thermal contributions from early (quark matter) and late (hadronic) phases dominate corresponding to small and large radial flow respectively.

In Fig.1, the invariant mass distribution of dileptons has been shown for a temperature $T = 0.2$ GeV. It is clear that the quark matter dominates over the hadronic matter contributions in the mass windows below the $\rho$-peak and above the $\phi$ - peak. It is expected that the slope parameters extracted from the transverse mass distribution of lepton pairs for mass region below the $\rho$ and above the $\phi$-peaks should be similar to quark matter phase. Therefore, slopes at these $M$ region will correspond to the early time when the radial follow is small. On the other hand the contributions in the region of $\rho$ mass is overwhelmingly from the hadronic phase and hence the slope at this region correspond to the late time containing larger radial flow.

First we discuss results for scenario (i) - here we have assumed that a thermalized state of quarks and gluons is formed after the collisions which reverts to hadronic phase through a very weak first order phase transition. The value of $\Gamma$ in eq. 9 is taken as 25 MeV. The values of initial temperature $T_i =$ 200 MeV, thermalization time $\tau_i = 1$ fm and the transition temperature $T_c = 175$ MeV have been assumed. We take the freeze-out temperature $T_f =$ 135 MeV which can reproduce the slope of the $\phi$ spectra measured by NA60 collaboration for In+In collisions [27]. With these inputs transverse mass spectra of dileptons have been evaluated for different invariant mass windows.

The resulting $m_T - M$ spectra of $\mu^+\mu^-$ are compared with the experimental data obtained by the NA60 collaboration for In-In collision for $\sqrt{s_{\rm NN}} =$ 17.3 GeV for different mass window in Figs. 2-5. The present calculation agrees well with the data which is shown as solid curves for all the mass ranges.

Finally in Fig 6 the effective temperature obtained from the inverse slope of these spectra have been plotted and compared with the data. The slopes have been estimated from theoretical results (shown by solid lines in Figs. 2-5) by parameterizing to an exponential function within the $(m_T - M)$ range $0.3 \leq m_T - M$(GeV)$\leq 1.0$. It is clear from the results that the slope at high $M$ region is reproduced well if the source at high $M$ region is predominantly partonic. A similar non-monotonic behaviour is observed in the variation of the elliptic flow ($v_2$) of photons as a of transverse momentum [28, 29].

Next we discuss results for scenario (ii). In this scenario we evolve a hadronic matter of initial temperature, $T_i$=200 MeV up to freeze-out temperature $T_f = 135$ MeV with an equation of state which includes mesons and baryons up to mass 2.5 GeV. The equation of state obtained for such a scenario is soft compared to an initial QGP state, the average velocity of sound ($c_s$) is $c_s^2 = 0.16$. With this scenario we fail to reproduce the data.

Keeping the values of $T_i = 200$ MeV and $T_f =$ 135 MeV fixed as above we increase the value of $T_c$ to 192 MeV and evaluated the slopes again for all the invariant mass windows shown in Figs. 2-5. In this case the contributions from the QGP phase is considerably reduced so the slopes of the $m_T$ spectra is pre-dominantly determined by the hadronic phase for all the invariant mass windows. The slopes for $M$ beyond the $\phi$ peak is not determined by the (early) QGP phase but by the (late) hadronic phase. As a



FIG. 6: The effective temperature obtained from the inverse slope of the invariant mass spectra. The solid line indicates results for $T_i = 200$ MeV, $T_c = 175$ MeV and $T_f = 135$ MeV.

result we observe a monotonically increasing trend of $T_{\rm eff}$ as a function of $M$. This also indicates that the source of lepton pairs at large ($M \sim 1.2$ GeV) is partonic. The sensitivity of $T_{\rm eff}$ to $T_c$ for various $M$ windows may be used to constrain the transition temperature.

In summary we find that the experimentally measured slope of the dilepton spectra by the NA60 collaboration can be explained theoretically if one assumes the formation of a partonic phase initially which reverts to hadronic phase through a weak first order or continuous transition. A strong first order transition does not reproduce the data. A hadronic initial state with realistic equation of state fails to reproduce the data. It is further observed that the spectra is sensitive to the value of the transition temperature. A larger value of $T_c$ (closer to $T_i$ in this case) fails to reproduce the non-monotonic behaviour of the slope, because in this case most of the contributions for all the invariant mass windows to the lepton spectra come from the late hadronic phase resulting in monotonic increase of $T_{\rm eff}$ with $M$. Therefore, the study of $T_{\rm eff}$ may be used to estimate $T_c$.

Currently the modification of hadronic spectral functions in a thermal bath and its effects on electromagnetic radiation is a field of great interest. The invariant mass distribution of lepton pairs are sensitive to both the pole shift and broadening [5, 6, 30, 31, 32, 33, 34, 35, 36, 37]. But the $p_T$ spectra of the EM radiation is insensitive to the broadening of the spectral function provided the integration over the $M$ is performed over the entire region. This is because broadening does not change the density of vector mesons significantly (see also [33]). However, the number density of vector mesons depends on the nature (shape) of the spectral function within the integration limit. Therefore, the $p_T$ spectra may change due to broadening when the integration over $M$ is done in a limited $M$ domain. By introducing the broadening of $\rho$ through a toy model $\Gamma_\rho = \Gamma^0_\rho(1 - T^2/T_c^2)^{-1}$ (where $\Gamma^0_\rho = 150$ MeV is the width of the $\rho$ in vacuum) we have observed that the resulting change in $T_{\rm eff}$ is small.

We are grateful to Sanja Damjanovic and Hans Specht for providing us the experimental data and having a fruitful discussion. We would like to thank Bedangadas Mohanty and Nu Xu for helpful discussions.

[1] L. D. McLerran and T. Toimela, Phys. Rev. D **31**, 545 (1985).
[2] C. Gale and J.I. Kapusta, Nucl. Phys. B **357**, 65 (1991).
[3] H.A. Weldon, Phys. Rev. D **42**, 2384 (1990).
[4] J. Alam, S. Raha and B. Sinha, Phys. Rep. **273**, 243 (1996).
[5] J. Alam, S. Sarkar, P. Roy, T. Hatsuda and B. Sinha, Ann. Phys. **286**, 159 (2000).
[6] R. Rapp and J. Wambach, Adv. Nucl. Phys. **25**, 1(2000).
[7] R. Arnaldi et al for the NA60 collaboration, Phys. Rev. Lett. **100** 022302 (2008).
[8] H van Hees and R. Rapp, Phys. Rev. Lett. **97** 102301 (2006).
[9] J. Ruppert, C. Gale, T. Renk, P. Lichard and J. I. Kapusta, Phys. Rev. Lett. **100** 162301 (2008).
[10] T. Renk and J. Ruppert, Phys. ReV. C **77**, 024907 (2008).
[11] K. Dusling and I. Zahed, hep-ph0701253.
[12] T. Altherr and P. V. Ruuskanen, Nucl. Phys. B **380**, 377 (1992).
[13] M. H. Thoma and C. T. Traxler, Phys. ReV. D **56**, 198(1997).
[14] E. V. Shuryak, Rev. Mod. Phys. **65** 1 (1993).
[15] P. Lichard and J. Juran, Phys. Rev. D **76** 094030 (2007).
[16] J. D. Bjorken, Phys. Rev. D **27**, 140 (1983).
[17] H. von Gersdorff, M. Kataja, L. D. McLerran and P. V. Ruskanen, Phys. Rev. D **34** 794(1986).
[18] R. C. Hwa and K. Kajantie, Phys. Rev. D **32**,1109 (1985).
[19] Y. Aoki, Z. Fodor, S. D. Katz and K. K. Szab, Phys. Lett. B **643** 46 (2006).
[20] M. Cheng et al., Phys. Rev. D **74**, 054507 (2006).
[21] B. Mohanty and J. Alam Phys. Rev. C **68** 064903 (2003).
[22] C. Bernard et al, Phys. Rev. D **75**, 094505 (2007).
[23] M. Asakawa and T. Hatsuda Phys. Rev. D **55** 4488 (1997).

[24] P. Braun-Munzinger, J. Stachel, J. P. Wessels and N. Xu, Phys. Lett. B **344** 43 (1995); **365** 1 (1996); P. Braun-Munzinger, I. Heppe and J. Stachel, *ibid* **465** 15 (1999); J, Cleymans and K. Redlich, Phys. Rev. Lett. **81** 5284 (1998); F. Becattini, J. Cleymans, A. Keranen, E. Suhonen and K. Redlich, Phys. Rev. C **64** 024901 (2001); P. Braun-Munzinger, D. Magestro, K. Redlich and J. Stachel, Phys. Lett. B **518** 41 (2001); M. Kaneta and N. Xu, J. Phys. G **27** 589 (2001).

[25] T. Hirano and K. Tsuda, Phys. Rev. C **66** 054905 (2002).

[26] E. Cooper and G. Frye, Phys. Rev. **10** 186 (1974).

[27] M. Floris for the NA60 collaboration, J. Phys. G**35** 104054 (2008).

[28] R. Chatterjee, E. S. Frodermann, U. W. Hienz and D. K. Srivastava, Phys. Rev. Lett. **96**, 202302 (2006).

[29] F.-M. Liu et al. (under preparation).

[30] G. E. Brown and M. Rho, Phys. Rev. Lett. **66** 2720 (1991).

[31] V.L. Eletsky, M. Belkacem, P.J. Ellis and J. I. Kapusta, Phys. Rev.C **64** 035202 (2001).

[32] M. Harada and C. Sasaki, Phys.Lett.B **537** 280 (2002).

[33] J. Alam, P. Roy, S. Sarkar and B. Sinha, Phys. Rev. C **67** 054901 (2003).

[34] S. Sarkar, J. Alam and T. Hatsuda, J. Phys. G **30** 607 (2004).

[35] G. Q. Li, C. M. Ko and G. E. Brown *Nucl. Phys.* A **606** 568 (1996).

[36] R. Rapp, G. Chanfray and J. Wambach, Nucl. Phys. A **617** 472 (1997).

[37] R. S. Hayano and T. Hatsuda arXiv:0812.1702 [nucl-ex]